# scientific reports

# OPEN



# Respiratory syncytial virus in pediatric patients with severe acute respiratory infections in Senegal: findings from the 2022 sentinel surveillance season

Mamadou Malado Jallow<sup>1</sup>, Moussa Moise Diagne<sup>1</sup>, Samba Niang Sagne<sup>2</sup>, Fatime Tall<sup>3</sup>, Jean Baptisse Niokhor Diouf<sup>4</sup>, Djibril Boiro<sup>5</sup>, Marie Pedapa Mendy<sup>1</sup>, Ndiendé Koba Ndiaye<sup>1</sup>, Davy Kiori<sup>1</sup>, Sara Sy<sup>1</sup>, Déborah Goudiaby<sup>1</sup>, Cheikh Loucoubar<sup>2</sup>, Gamou Fall<sup>1</sup>, Mamadou Aliou Barry<sup>2</sup> & Ndongo Dia<sup>1⊠</sup>

In 2022, many regions around the world experienced a severe respiratory syncytial virus (RSV) epidemic with an earlier-than-usual start and increased numbers of paediatric patients in emergency departments. Here we carried out this study to describe the epidemiology and genetic characteristics of RSV infection in patients hospitalized with severe acute respiratory infections in 2022. Samples were tested for RSV by multiplex real time reverse transcription polymerase chain reaction. Subsequently, a subset of RSV positive samples was selected for NGS sequencing. RSV was detected in 16.04%, among which RSV-A was confirmed in 7.5% and RSV-B in 76.7%. RSV infection were more identified in infants aged ≤11 months (83.3%) and a shift in the circulation pattern was observed, with highest incidences between September–November. Phylogenetic analyses revealed that all RSV-A strains belonged to GA2.3.5 genotype and all RSV-B strains to GB5.0.5a genotype. Three putative N-glycosylation sites at amino acid positions 103, 135, 237 were predicted among RSV-A strains, while four N-linked glycosylation sites at positions 81, 86, 231 and 294 were identified in RSV-B strains. Globally, our findings reveal an exclusive co-circulation of two genetic lineages of RSV within the pediatric population in Senegal, especially in infants aged ≤11 months.

Acute lower respiratory infection (ALRI) caused by respiratory syncytial virus (RSV) has gained recognition as a global health problem with a high burden of disease<sup>1</sup>. Infants and children under 5 years old are particularly susceptible to severe diseases caused by RSV, manifesting as a spectrum of upper and/or lower respiratory tract infections including bronchiolitis and pneumonia<sup>2</sup>. Globally, the virus has led to 33 million episodes of ALRI, 3.6 million hospital admissions, 26,300 in-hospital deaths, and 101,400 RSV-attributable overall deaths in children younger than 5 years<sup>3</sup>. In addition, 99% of RSV-related deaths globally occur in low- and middle-income countries, making such infections a high priority for this setting<sup>4</sup>. Although often characterized as a pediatric disease, RSV infection in adults represents a substantial health burden<sup>5</sup>. Mortality attributable to RSV in adults aged 65 years or older is estimated to be 7.2 per 100,000 person-years, and 8% of RSV ALRI among older adults admitted to hospital was reported to result in death in the USA<sup>1</sup>. RSV is highly seasonal as RSV epidemics tend to occur in the winter in temperate regions<sup>6</sup> and during the rainy season in tropical countries<sup>7</sup>. Despite the significant public health impact and global economic burden imputed to this pathogen, approved vaccines were recent<sup>8</sup> with a currently limited access, especially for low income resources countries were RSV immunization program are lacking. Also, a RSV monoclonal antibody product (palivizumab) is available for immunoprophylaxis in a limited high-risk infant population<sup>9</sup>. Therefore, for the time being, measures preventing the spread of RSV remain the most promising means of controlling these seasonal epidemics<sup>10</sup>.

<sup>1</sup>Département de Virologie, Institut Pasteur de Dakar, Dakar, Senegal. <sup>2</sup>Institut Pasteur de Dakar, Unité d'Epidémiologie Des Maladies Infectieuses, 36, Avenue Pasteur, B.P. 220, Dakar, Senegal. <sup>3</sup>Hôpital Des Enfants Albert Royer de Fann, Dakar, Senegal. <sup>4</sup>Hôpital Roi Baudoin de Guediawaye, Dakar, Senegal. <sup>5</sup>Centre Hospitalier Abass Ndao, Dakar, Senegal. <sup>⊠</sup>email: ndia@pasteur.sn

Respiratory syncytial virus (RSV) is a non-segmented negative-sense single-strand ribonucleic acid (RNA) genome of approximately 15.2 kb packaged in a lipid envelope<sup>2,11</sup>. This genome encodes eleven proteins, including attachment (G) and fusion (F) glycoproteins which contain neutralising antibody epitopes capable of inducing a neutralising antibody response, and are targeted in RSV vaccine development strategies<sup>12</sup>. This enveloped virus is a member of the genus *Ortopneumovirus* in the family *Pneumoviridae* and has only one serotype, divided into antigenic subgroups A (RSV-A) and B (RSV-B)<sup>13,14</sup>. Within each subgroup there are numerous genotypes, which have historically been classified based on the gene sequences of the Hypervariable Region 2 (HVR2) of glycoprotein G<sup>15,16</sup>. Usually, both subtypes co-circulate during seasonal epidemic periods with alternating patterns of predominance over time<sup>17</sup>, and currently the most frequent genotypes worldwide are ON1 for RSV-A and BA for RSVB, characterized by duplications of a 72 and a 60 nucleotides in the G gene, respectively<sup>18,19</sup>.

In December 2019, the world experienced the beginning of the coronavirus disease 2019 (COVID-19) pandemic, caused by a novel coronavirus designated as severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2)<sup>20</sup>. The global impact at the public health level was catastrophic, with millions of hospitalizations and deaths<sup>17</sup>. Senegal confirmed its first COVID-19 case on March 2, 2020 when the country had already prepared for its arrival, including rapid detection of cases, patient's isolation, tracing and quarantine of contacts<sup>21</sup>. Nevertheless, COVID-19 cases increased, and consequently mitigation measures were taken by health authorities, including travel restrictions and border closures, curfews, physical distancing and mandatory wearing of face masks as part of a comprehensive lockdown<sup>22</sup>. These sanitary measures adopted to reduce SARS-CoV-2 transmission, not only impacted COVID-19 but may have also deeply modified the natural course of seasonal viral infections, such as RSV<sup>23,24</sup>. However, with the easing of COVID-19 imposed restrictions, many regions experienced a severe RSV epidemic with an earlier-than-usual start and increased numbers of pediatric patients in emergency departments, partly due to a lack of protective immunity in the community following a lack of exposure from the previous season<sup>10,25,26</sup>. In Senegal, a surveillance of Severe Acute Respiratory Infections (SARI) which falls within the scope of the 4S (Sentinel Syndromic Surveillance in Senegal) network activities was set up in healthcare centers in the capital city Dakar since 2015 with the aim to allow the Ministry of Health to quickly detect and alert any abnormal health event<sup>27</sup>. So here, we carried out a study which describes the epidemiology and genetic characteristics of RSV infection in patients hospitalized with SARI in 2022 after the alleviation of non-pharmaceutical interventions due to the COVID-19 pandemic.

# Results

### **RSV** confirmed cases

During January, 1 to December 31 2022, a total of 748 SARI samples were received at the National Influenza Center and analyzed by multiplex reverse transcription polymerase chain reaction (RT-PCR). Of these patients, 52.3% (391/748) were male and the median age of the study population was 11 months (interquartile range [IQR]: 4.8 years), with 50.7% (379/748) of all patients being  $\leq$  11 months old. Overall, RSV was detected in 16% of all samples (120/748), among which RSV type A (RSV-A) was confirmed in 10% (12/120), whereas RSV type B (RSV-B) was encountered in 89.2% (107/120). For the remaining RSV positive sample (0.8%; 1/120), the type could not be determined due to a low viral load. (Table 1). Coinfections of RSV-A and RSV-B were not encountered in any of tested samples. RSV-B cases outnumbered those of the RSV-A throughout the study period as showed in Fig. 1. During the first trimester of this study (calendar week 1 to calendar week 14), there was almost no circulation of RSV with only 1 case (RSV-A) in the first calendar week of the year (January). However, we noticed a gradual increase in RSV cases from calendar week 24, reaching a peak in activity between weeks 38 and 40 (September–October) (Fig. 1).

#### Characteristics of inpatients infected with RSV

Demographic characteristics and clinical parameters of RSV-associated SARI patients are presented in Table 1. Among the 120 RSV-positive patients, 55 were male and 65 were female. Their age ranged from 1 month to 31 years, with mean and median ages of 10 months (0.84 year) and 2.4 months (0.2 year) (interquartile range [IQR]: 0.59 year) respectively. With regards to the distribution by age group, we noted that infants aged  $\leq 11$  months were the most infected group with a detection rate of 83.3% (100/120). RSV infection rapidly dropped after the age of 1 year, with detection rates decreasing from 9.2% in patients aged 1–2 years old (11/120) to 0% in patients aged 11–15 years old (0/120). Only one case of RSV (0.8%) was encountered in patients > 15 years old. At the time of admission, besides cough (75.8%) and fever (43.3%) which were part of case-definition, we found that breathing difficulties (adjusted OR 2.02; 95% CI 1.27–3.37) were more commonly reported among RSV-associated SARI patients (*P-value* = 0.004). Taking into account the clinical diagnosis, more than half of all inpatients infected with RSV had bronchiolitis (57.5%), whereas pneumonia was diagnosed in 37 patients (30.8%). Asthma exacerbation and acute bronchitis were diagnosed in 11 (9.2%) and 8 (6.7%) RSV-associated SARI patients respectively. Antibiotics and bronchodilators were administrated to 65.0% (78/120) and 33.3% (40/120) of patients, respectively (Table 1).

#### Co-infection of RSV with other respiratory pathogens

Co-infections among the 120 RSV-positive cases with other community-acquired respiratory pathogens were encountered in 56 (46.7%) patients. Mixed infections with respiratory bacteria were the most commonly observed in this study, with *Haemophilus Influenzae* (17.5%) and *Streptococcus Pneumoniae* (14.2%) being the most co-detected pathogens. Co-infection with *Klebsiella Pneumoniae* was observed in 8 patients (6.7%), *Moraxella Pneumoniae* in 5 patients (4.2%), *Staphylococcus aureus* in 3 patients (2.5%) and *Bordetella Pertussis* (bacterium responsible for whooping cough) in a single patient (0.8%). The most common co-infecting viruses were

		Total tested (n=748)						
RSV positive	Overall	RSV A	RSV B	Untyped RSV	p-value	Crude OR	Adjusted OR	
Characteristics	(n=120)	(n=12)	(n=107)	(n=1)		(95% CI)	(95% CI)	
Gender, n (%)								
Male	55 (45.8)	6 (50.0)	49 (45.8)	0 (0.0)	1	1	1	
Female	65 (54.2)	6 (50.0)	58 (54.2)	1 (100)	0.1	0.73 [0.5—1.1]	0.69 [0.45-1.05]	
Age group, n (%)								
≤11 months	100 (83.3)	11 (91.7)	88 (82.2)	1 (100)	1	1	1	
1-2 years	11 (9.2)	1 (8.3)	10 (9.3)	0 (0.0)	5.63 e <sup>-05</sup>	0.3 [0.1-0.5]	0.28 [0.15-0.5]	
3–5 years	5 (4.2)	0 (0.0)	5 (4.7)	0 (0.0)	0.001	0.1 [0.03-0.4]	0.15 [0.036-0.4]	
6-10 years	2 (1.7)	0 (0,0)	2 (1.9)	0 (0,0)	0.007	0.07 [0.004-0.3]	0.06 [0.003-0.3]	
11-15 years	0 (0.0)	0 (0,0)	0 (0.0)	0 (0.0)	0.9	-	-	
>15	1 (0.8)	0 (0,0)	1 (0.9)	0 (0,0)	0.001	0.04 [0.002-0.17]	0.03 [0.002-0.16]	
Mean age (95% CI)	0.8 [0.3-1.4]	0.2 [0-0.3]	0.9 [0.3–1.5]	-		-	-	
Clinical manifestation, n (%)								
Fever	52 (43.3)	12 (100)	107 (100)	1 (100)	0.0003	0.48 [0.32-0.72]	0.46 [0.31-0.69]	
Cough	91 (75.8)	12 (100)	107 (100)	1 (100)	0.62	1.12 [0.72-1.78]	1.12 [0.71—1.79]	
Breathing difficulties	97 (80.8)	11 (91.7)	85 (79.4)	1 (100)	0.004	2.01 [1.26-3.33]	2.02 [1.27-3.37]	
Rhinitis	6 (5.0)	0 (0,0)	5 (4.7)	0 (0.0)	0.96	0.98 [0.36-2.24]	0.95 [0.35-2.19]	
Vomiting	2 (1.2)	0 (0,0)	2 (1.9)	0 (0.0)	0.51	0.61 [0.09-2.16]	0.61 [0.09-2.16]	
LRT complications, n (%)								
Pneumonia	37 (30.8)	3 (25.0)	33 (30.8)	1 (100)	<2.2e <sup>-16</sup>	-	-	
Bronchiolitis	69 (57.5)	8 (66.7)	61 (57.0)	0 (0.0)	< 2.2e <sup>-16</sup>	-	-	
Acute bronchitis	8 (6.7)	1 (8.3)	7 (6.5)	0 (0.0)	1.73e <sup>-09</sup>	-	-	
Asthma exacerbation	11 (9.2)	1 (8.3)	10 (9.3)	0 (0.0)	4.83e <sup>-13</sup>	-	-	
Prescribed treatment, n (%)								
Antibiotics	78 (65.0)	10 (83.3)	67 (62.6)	1 (100)	< 2.2e <sup>-16</sup>	-	-	
Bronchodilators	40 (33.3)	3 (25.0)	37 (34.6)	0 (0.0)	< 2.2e <sup>-16</sup>	-	-	

 Table 1. Demographic characteristics and clinical parameters of RSV-associated SARI patients, Senegal

 January-December 2022. n number of patients, LRT lower respiratory tract, RSV respiratory syncytial virus.



**Figure 1.** Weekly distribution of RSV among inpatients in Senegal from January to December 2022. Bars represent the proportions of RSV cases for each epidemiological week and the curve represent the positivity rates.

	RSV A	RSV B	Untyped RSV	Overall		
Pathogens	(n=12)	(n=107)	(n=1)	(n=120)		
Viruses, n (%)						
SARS-CoV2	0 (0.0)	5 (4.7)	0 (0.0)	5 (4.2)		
Adenovirus	0 (0.0)	2 (1.9)	0 (0.0)	2 (1.7)		
HCoV-OC43	0 (0.0)	3 (2.8)	0 (0.0)	3 (2.5)		
Enterovirus	0 (0.0)	1 (0.9)	0 (0.0)	1 (0.8)		
HMPV	0 (0.0)	2 (1.9)	0 (0.0)	2 (1.7)		
Parainfluenza virus	0 (0.0)	2 (1.9)	0 (0.0)	2 (1.7)		
Rhinovirus	2 (16.7)	8 (7.5)	0 (0.0)	10 (8.3)		
Bacteria, n (%)						
Haemophilus influenza	2 (16.7)	19 (17.7)	0 (0.0)	21 (17.5)		
Klebsiella pneumoniae	0 (0.0)	7 (6.5)	1 (100)	8 (6.7)		
Bordetella Pertussis	0 (0.0)	1 (0.9)	0 (0.0)	1 (0.8)		
Staphylococcus aureus	1 (8.3)	1 (0.9)	1 (100)	3 (2.5)		
Moraxella pneumoniae	2 (16.7)	3 (2.8)	0 (0.0)	5 (4.2)		
Streptococcus pneumoniae	1 (8.3)	15 (14.0)	1 (100)	17 (14.2)		

**Table 2.** Co-infection of RSV with other respiratory pathogens. SARS-CoV2 severe acute respiratorysyndrome coronavirus 2, HCoV-OC43 human Coronavirus OC43, HMPV human metapneumovirus.

rhinovirus accounting for 10 cases (8.3%), followed by SARS-CoV2 with 5 cases (4.2%), and HCoV-OC43 with 3 cases (2.5%). RSV/HMPV co-infections were detected in 2 RSV-associated SARI patients (1.7%) (Table 2).

### Phylogenetic analysis of RSV-A and RSV-B sequences

Overall, 7 RSV-A and 29 RSV-B whole genomes were successfully obtained in this study. The Nextclade online tool (https://clades.nextstrain.org/, accessed 2 August 2023) were initially used for genotype predictions according to Goya et al. 2019<sup>28</sup>. Genotyping of the G gene found that all RSV-A sequences were GA2.3.5 lineage (clade A23), and all RSV-B sequences were GB5.05a lineage (clade B6). We undertook a more detailed phylogenetic analysis of complete genomes with historical and recent RSV sequences in public databases (Genebank and GISAID) up to December 2022 to ascertain more precisely the genetic relationships among RSV strains generated as part of this study and their putative closest ancestors. Phylogenetic trees showed that Senegalese strains were closely related to RSV strains circulating globally in the same period. All RSV-A sequences obtained in this study belonged to the GA2.3.5 genotype (ON1 strains) (Fig. 2), and all RSV-B sequences were classified as genotype GB5.0.5a (BA9 strains) (Fig. 3).

# Genetic analysis of deduced amino acid sequences of RSV strains from Senegal

Deduced amino acid sequences of the G protein of RSV-A and RSV-B strains from Senegal were aligned and compared with prototype strain A2 (GenBank accession number M11486) and ON1 reference strain (GenBank accession number JN257693) for RSV-A and BA4128/99B strain (GenBank accession number AY333364) for RSV-B. For Senegalese RSV-A strains, the 72 nucleotides duplication in the C-terminal end of the G gene, which is characteristic to ON1 strains<sup>18</sup> was observed in five sequences, leading to an insertion of 24 amino acid (QEETL-HSTTSEGYLSPSQVYTTSG) of which 23 are duplications of amino acid spanning positions 261-283 (Fig. 4A). Analysis of all viral genes showed several amino acid substitutions compared to hRSV/A/England/397/2017 (EPI\_ISL\_412866). All RSV-A samples from Senegal shared mutations P71L, H90Y, I134K, G224E and S243I in the G attachment glycoprotein, V352A in the N protein, L55P in the P protein and S176P in the M2-1 protein. With regards to the fusion glycoprotein F, mutations A103T and T122A which is believed to remove a potential N-glycosylation site at amino acid position 120 (the motif at positions 120-122 changed from NNT (glyco) to NNA (no glyco) were observed only in strains with the 24 amino acid insertion. Using the NetNGlyc 1.0 server, all RSV-A sequences were predicted to have three putative N-glycosylation sites at amino acid positions 103-106 (NLSG), 135–138 (NTTT) and 237–240 (NTTK) such as the ON1 reference strain (JN257693), with the exception of two sequences (hRSV/A/Senegal/1017/2022 and hRSV/A/Senegal/2037/2022) with the substitution T135I leading to the loss of one N-glycosylation site. When compared to the RSV-A prototype strain A2, only one of the three potential N-glycosylation sites predicted in this study remains conserved between all Senegalese RSV-A isolates (amino acid 237 in RSV-A prototype strain A2).

With regards to RSV-B strains, the presence of 60 nucleotides duplication in the 2nd HVR of the G gene, which is a characteristic of the BA genotypes<sup>19</sup>, was found in all sequences, resulting in a duplication of 20 amino acids (TERDTSTSQSTVLDTTTSKH) spanning positions 238–257 (Fig. 4B). In comparison with the hRSV/B/Australia/VIC-RCH056/2019 reference strain (EPI\_ISL\_1653999) using the GISAID RSVsurver, several mutations were found in most of the viral genes, including the G gene where several amino acid substitutions were found, of which A74V, T131A, I137T, I252T and I268T were present in all RSV-B sequences obtained in this study. All RSV-B strains shared the V97I mutation in the N protein and T1987I in the L protein. Substitutions in the fusion glycoprotein, such as R191K, M206I, R209Q, P312H, S190N, S211N and S389P were identified in



0.03

**Figure 2.** Phylogenetic tree of complete genomes of RSV-A subtype circulating among hospitalized patients in Senegal (January–December 2022). The tree was constructed using the maximum likelihood (ML) method using the IQ-TREE software version 1.6.12<sup>18</sup> and visualized using the Figtree software version 1.4.4. The statistical significance was tested by 1000 bootstrapping replicates, and the software was responsible for defining the correct model used. Sequences from Senegal are highlighted in red color. The scale bar represents the number of nucleotide substitutions per site.

most RSV-B sequences from Senegal. When compared with the BA4128/99B strain, S245P and V269A amino acid substitutions in the duplicated 20 amino acid region were observed in all RSV-B strains. The analysis of potential N-glycosylation sites revealed four conserved N-linked glycosylation sites at positions 81–84 (NHTE), 86–89 (NITT), 228–231 (NPTK) and 294–297 (NSTQ) in the 2nd HVR of the G gene of all RSV-B strains in this study in comparison with the BA4128/99B prototype reference strain. The amino acid substitution T310I resulted in a loss of a potential N-glycosylation site due to the mutation K256N located in the duplicated 20 amino acid region. Furthermore, amino acid substitution N6S observed in one sample (hRSV/B/Senegal/2031/2022) resulted in the acquisition of a potential N-linked glycosylation.

# Discussion

Before the COVID-19 pandemic, respiratory syncytial virus (RSV) usually circulated primarily in the second half of each year, between June and September in Senegal, which coincide with the period of the rainy season<sup>29</sup>. However, the activity of this virus was disrupted during the COVID-19 pandemic, which circulated at historically low levels due to implementation of public health measures to prevent the spread of SARS-CoV-2 and mitigate the impact of the pandemic. In 2022, after the alleviation of COVID-19 measures, Senegal, like several other countries<sup>26,30–32</sup>, noticed an unprecedented number of RSV detections with a temporal shift (peak of detection between September and October) and increased numbers of pediatric inpatients in the different hospital sentinel sites of the 4S network. The reasons for this atypical resurgence of RSV cases are unclear, but it may be due to a deficit in RSV immunity due to prolonged lack of viral exposure<sup>33</sup>. Furthermore, this resurgence can be



0.005

**Figure 3.** Phylogenetic tree of complete genomes of RSV-B subtype circulating among hospitalized patients in Senegal (January-December 2022). The tree was constructed using the maximum likelihood (ML) method using the IQ-TREE software version 1.6.12<sup>18</sup> and visualized using the Figtree software version 1.4.4. The statistical significance was tested by 1000 bootstrapping replicates, and the software was responsible for defining the correct model used. Sequences from Senegal are highlighted in red color. The scale bar represents the number of nucleotide substitutions per site.

explained by specific changes in the circulating genotypes and/or the emergence of a novel strains with increased transmission or pathogenicity<sup>26</sup>. Therefore, to understand whether the increased number of RSV cases seen in the post-pandemic period, was due to specific changes in the circulating genotypes or to the emergence of novel strains, and whether they cause severe illness, we investigated the epidemiology and the genetic characteristics of RSV circulating among hospitalized patients with SARI in Senegal from January to December 2022. Among the 748 samples from the hospital-based surveillance screened as part of this study, 120 were positive for RSV, representing a detection rate of 16.04%. This prevalence is higher than the rate of 11.4% reported in a previous study in Senegal in Influenza-like Illness (ILI) outpatients after four consecutive years of surveillance (2012-2015)<sup>29</sup>. However, many other similar studies reported higher detection rates of RSV infection, including Thailand<sup>34</sup> with 27.2%, Germany<sup>35</sup> with 42.8%, Italy<sup>36</sup> with 40.6% and Bulgaria<sup>37</sup> with 26.2%. On the other hand, and in accordance with our current findings, a report in China indicated an overall positivity rate of 16.0% (95% CI=12.9, 19.6%) among children<sup>38</sup>. The possible reasons for the difference in RSV infection rates could be reflecting the true burden in each region, or it is possibly attributed to the use of different diagnostic methods, study populations (outpatients or hospitalized patients), the sampling period, environmental factors and even the duration of the study. As reported in various countries<sup>26,30,39</sup>, the 2022 surge in Senegal was mainly caused by the RSV-B subtype, with 89.2% of the overall RSV positive cases. Unlike our results, several authors reported RSV-A as the dominant subtype<sup>35,40</sup>. Our study found no mixed infection of RSV-A/RSV-B, despite the reporting of 3.4% of RSV-A/B coinfection in ILI patients in a previous study in Senegal by Fall et al.<sup>29</sup> or the 2% reported by Hall et al.<sup>41</sup> in a population-based surveillance of acute respiratory infections among children under 5 years of age in three U.S. counties. However, our findings regarding Haemophilus Influenzae and Streptococcus Pneumoniae as the most frequently co-detected bacteria with RSV were similar to those of Lin et  $al^{42}$ . According to some studies, coinfections can exacerbate the severity of RSV disease while others found no conclusive evidence for a A

SV-A prototype strain A2 hSV/A/Senegal/2255/2022 hSV/A/Senegal/1712/2022 hSV/A/Senegal/0653/2022 bSV/A/Senegal/0653/2022	100         110         120         130         140         230         240         250         260         270         280         290         300         310         320           100         110         120         130         140         230         240         250         260         270         280         290         300         310         320           100         110         120         130         140         250         260         270         280         290         300         310         320           100         100         130         140         250         260         270         280         290         300         310         320           100         100         150         160         170         150         160         170         180
hRSV/A/Senegal/0748/2022 hRSV/A/Senegal/1017/2022 hRSV/A/Senegal/2037/2022	.F.L.GT.S
ON67-1210A strain	.F. L.GTS
B	70 80 - 50 100 110 120 130 140 200 210 220 230 240 250 260 270 280 250 350 350 350 350 350 350 350 350 350 3
hRSV/B/Senegal/2358/2022	
hRSV/E/Sepegal/2359/2022	
hRSV/B/Senegal/0850/2022	
hESV/B/Senegal/1022/2022	1
hRSV/B/Senegal/1847/2022	
hESV/B/Senegal/1244/2022	
hRSV/B/Senegal/1103/2022	1
hR3V/B/Senegal/1195/2022	
hRSV/B/Senegal/1344/2022	1
hR3V/B/Senega1/1345/2022	han a second s
hR3V/8/Senegal/1478/2022	and a second s
hR8V/B/Senegal/1955/2022	
hR5V/B/Senegal/1956/2022	and a second
hR3V/B/3enegal/2162/2022	
hRSV/B/Senegal/2170/2022	
hRSV/B/Senegal/2258/2022	
hR3V/B/3enegal/2255/2022	
hRSV/B/Senegal/2293/2022	
hR5V/B/Senegal/2456/2022	
hR3V/B/Senegal/2547/2022	
hRSV/B/Senegal/0486/2022	
hRSV/B/Senegal/0847/2022	
hRSV/B/Senegal/1014/2022	
nBUV/B/Benegal/0846/2022	
hRSV/B/Senegal/1631/2022	N N N N N N N N N N N N N N N N N N N
hRSV/B/Senegal/1640/2022	
BEEV/B/Senegal/2353/2022	
The second state of second state of the second	
hRSV/8/Senegal/2031/2022	

**Figure 4.** Deduced amino acid alignment and mutations in the second hypervariable region of the G protein of RSV-A (**A**) and RSV-B (**B**) strains from Senegal compared with prototype strain A2 (GenBank accession number M11486) and ON1 reference strain (GenBank accession number JN257693) for RSV-A, BA4128/99B strain (GenBank accession number AY333364) for RSV-B. The two copies of 23 amino acid regions in the RSV-A strains (**A**) and 20 amino acid regions in the RSV-B strains (**B**) are framed black, whereas potential N-glycosylation sites (NXT, where X is not proline) are indicated by red boxed areas. Identical amino acids are indicated by dot and asterisks represent stop codons.

link between the presence of coinfection and the disease severity<sup>43,44</sup>. Therefore, further studies will be needed to elucidate the relationships between RSV co-infection and disease severity.

Although this epidemic is atypical, it seems to have a typical age-related RSV risk pattern, with infants aged  $\leq 11$  months being the most affected groups (83.3% of the overall RSV positive cases) and the risk of RSV infection decreases with increasing age (9.2% in patients aged 1–2 years old and 0% in patients aged 11–15 years old) due to the development of immunity after repeated infections<sup>37</sup>. This reinforces the idea that the implementation of any prophylaxis (vaccine, monoclonal antibodies) should target children under 1 year. Consistent with our findings, several studies have reported this sensitivity of infants, especially in the first year life to RSV infection<sup>35,45–47</sup>. The relative immaturity of the immune system for most RSV genotypes in early childhood may explain this high prevalence of RSV infection in infant<sup>30</sup>. In line with several reports<sup>37,40</sup>, the majority of RSV-associated patients in this study presented with bronchiolitis (57.5%) followed by pneumonia (30.8%). Indeed, bronchiolitis and pneumonia are among the most important RSV-related concerns, as RSV can directly cause viral pneumonia or patients may be additionally co-infected with bacteria as shown in this study, and both can often lead to hospitalization and severe disease course, increasing the risk of ICU transfer, especially in at-risk groups<sup>42,46,48,49</sup>.

Phylogenetic analysis of all available complete genomes of RSV strains infecting patients hospitalized with SARI in Dakar, revealed an exclusive co-circulation of two genetic lineages, GA2.3.5 for RSV-A and GB5.0.5a for RSV-B within the population; however, GB5.0.5a strains dominated over GA2.3.5 during the entire study period. A similar genotypic composition has been observed in several other countries around the world<sup>26,47,50</sup>. These genotypes driving the 2022 RSV outbreak in Senegal had been circulating within the Senegalese population in pre-pandemic years<sup>14</sup>, suggesting that there are no specific changes in RSV since the COVID-19 pandemic began that would account for increased viral spread.

In agreement with previous findings around the world<sup>15,37,40</sup>, the genetic analysis of deduced amino acid indicated that Senegalese strains possessed several amino acid substitutions, particularly in the second hyper variable region of the G protein gene. Amino acid mutations, P71L, H90Y, I134K, G224E, S243I in RSV-A and A74V, T131A, I137T, I252T, I268T in RSV-B were detected in all sequences compared with reference strains hRSV/A/England/397/2017 and hRSV/B/Australia/VIC-RCH056/2019 respectively, which confirms the high genetic variability of the RSV G gene. Unlike the attachment G gene, few key amino acid changes were found in the other remaining genes for both subtypes, including V352A in the N protein, L55P in the P protein and S176P in the M2-1 protein of all RSV-A strains. Similar to the findings of Goya et al.<sup>50</sup>, the amino acid constellation A103T and T122A in the RSV-A fusion protein was encountered only in sequences with the 72 nucleotide duplication in the C-terminal end of the G gene. With regards to RSV-B strains, several mutations were found in most sequences, such as R191K, M206I, R209Q, P312H, S190N, S211N and S389P in the fusion glycoprotein F, V97I in the N protein and T1987I in the L protein. The virus evasion from host immune response recognition may be mediated by the presence of N- and O-linked glycans on the G protein. It has been established that N and O linked glycosylation in G protein can alter the attachment and antigenicity properties of RSV<sup>16,51</sup>. Unlike

the findings of Chen et al.<sup>51</sup> in China, three putative N-glycosylation sites at amino acid positions 103, 135, 237 were predicted among RSV-A strains, while four conserved N-linked glycosylation sites at positions 81, 86, 231 and 294 were identified in RSV-B strains.

However, we pointed out some limitations in our study. The major limitation of the present study is the fact that only hospitalized patients in Dakar were included, and thus our data may not represent the national RSV burden. Therefore, it would be important to extend the hospital-based surveillance in other regions to firmly establish the burden associated with RSV in Senegal, and clear correlations between disease severity and genotypes.

In summary, this hospital-based surveillance revealed that the abnormal increase in RSV-positive rates after the alleviation of the COVID-19 measures in Senegal, was mainly due to RSV-B subtype. Globally, we observed a shift in the circulation pattern of RSV (systematic increase in infections between August and November), which usually circulates during the rainy season between June through September each year. Phylogenetic analyses highlighted that all RSV-B strains belonged to GB5.0.5a (BA9 lineage) genotype very similar to that previously circulating, suggesting that the increased number of RSV cases, especially in infants aged  $\leq 11$  months is likely because of diminished protective immunity in the population from low RSV exposure, a consequence of pandemic mitigation measures. Therefore, a deep molecular surveillance of respiratory-associated viruses should be established under the COVID-19 strategy to avoid unexpected outbreaks of other diseases.

# Material and methods

#### Study design, sample and data collection

This study was conducted from January to December 2022 and samples were collected as part of the routine hospital-based surveillance of SARI in Senegal piloted by the National Influenza Centre (NIC) hosted at the Institut Pasteur de Dakar (IPD). A standardized SARI case definition according to the WHO was followed, which defines SARI as "an acute respiratory infection with history of fever or measured fever of  $\geq$  38 °C, and cough with onset within the last ten days and requires hospitalization"52. Participants were enrolled in seven sentinel hospitals (all located in the capital city Dakar) for the surveillance of SARI: Albert Royer Children's Hospital, Diamniadio Children's Hospital, General Hospital Idrissa Pouye of Grand Yoff, Fann Teaching Hospital, Principal Hospital of Dakar, Abass Ndao Hospital and Roi Baudouin Hospital (Fig. 5). Upon admission, nasopharyngeal and/or oropharyngeal swabs were collected per patient by qualified medical personnel. The swabs were inserted in a universal viral transport medium (Becton Dickinson and company, Milano, Italy) and promptly transported at a controlled temperature (4-8 °C) to the laboratory for the screening of respiratory pathogens, including RSV within the 24 h following samples collection. In addition, for each identified patient admitted at the selected hospitals, demographic, clinical, and epidemiological information for all SARI cases were recorded in a casebased surveillance form by trained nurses. Upon receipt in the laboratory, samples were tested using a multiplex real-time reverse transcription polymerase chain reaction assay and left-over clinical specimens were stored at - 80 °C for biobanking and additional analyses (e.g., NGS sequencing).

#### Nucleic acid extraction and screening of respiratory pathogens

Total nucleic acid was extracted from 200  $\mu$ l of swab suspension in Viral Transport Medium (VTM) using the QIAamp Viral RNA kit (QIAGEN, Valencia, CA, USA), according to the manufacturer's instructions. The



**Figure 5.** Map of Senegal showing the different sentinel sites of the 4S network. Hospital sentinel sites, all located in the capital Dakar are represented by red dots, while blue dots represent the ILI community sentinel sites.

extracted nucleic acid was eluted in a final volume of 60 µl and immediately used for the routine testing of respiratory pathogens. Extracted RNA from all samples was tested for RSV, along with other respiratory pathogens (Virus and bacteria) by multiplex real time reverse transcription polymerase chain reaction (rRT-PCR) system, using the Allplex™ Respiratory Full Panel Assay (Seegene, Seoul, Republic of Korea). This multiplex one-step realtime RT-PCR assay that composed of 4 different panels enables the simultaneous detection and differentiation of 26 causative pathogens in respiratory tract infections including 16 viruses, 3 Flu A subtypes and 7 bacteria. Respiratory syncytial virus A (RSV-A) and Respiratory syncytial virus B (RSV-B) along with the 3 Flu A subtypes (A/H1pdm09, A/H3 and A/H1) were detected with the Panel 1. Briefly, the total reaction volume for each real time PCR was 25 µl, consisting of 5 µl Nuclease-free water, 5 µl of 5XRP MOM (MuDT Oligo Mix), 5 µl of 5X Buffer (Buffer containing dNTPs), 2 µl of Enzyme Mix and 8 µl of each extracted or controls nucleic acid under the following cycling conditions: reverse transcription step of 20 min at 50 °C, initial denaturation step of 15 min at 95 °C, followed by 45 PCR cycles of 10 s at 95 °C, 1 min at 60 °C and 10 s at 72 °C. Fluorescence is detected at 60 °C and 72 °C. The real-time RT-PCR was done on the CFX96" (Bio-Rad, California, USA) platform, and subsequently interpreted by Seegene's Viewer software. The latter considers respiratory targets that generate adequate, exponential fluorescence curve with Ct-values or cycle threshold value (it represents the number of cycles required for the fluorescent signal to cross the threshold. It is inversely proportional to the amount of target nucleic acid in the sample), below 42 cycles as positive.

# Next generation sequencing of RSV positive samples

For RSV whole genome sequencing, stored positive samples with a relatively high viral load (Ct-value < 30) were selected and transmitted to the IPD sequencing platform to perform hybridization capture-based, metagenomic next-generation sequencing using the Twist Respiratory Virus Research Panel (103067; Twist Biosciences, San Francisco, CA), as previously described<sup>53</sup>. The panel is a culture-free, hybrid capture enrichment workflow embedded as a sub analysis panel in One Codex analysis platform and is capable of detecting 29 common human respiratory viruses simultaneously<sup>54</sup>. To generate the consensus genomes of RSV in all samples, raw reads from Illumina sequencing in FASTQ format were assembled by de novo assembling using CZ ID (https://czid.org/, accessed on June 21, 2023), a cloud based open-source bioinformatics pipeline for metagenomic sequencing data.

### Sequence alignments and phylogenetic analysis

Phylogenetic analyses were carried out by adding sequences from other countries strains downloaded from Gen-Bank (https://www.ncbi.nlm.nih.go, accessed on 21 June 2023) and GISAID (https://www.gisaid.org, accessed on 21 June 2023) databases, as well as adding reference sequences of known genotypes. Sequences alignments were performed by using MAFFT software implementing the FFT-NS-2 algorithm<sup>55</sup>. The alignment was manually inspected in Bioedit v7.1.3.0<sup>56</sup> to ensure accuracy and that identical sequences were removed. The alignment was used to construct phylogenetic trees by means of the maximum likelihood method by using IQ-TREE v.2.1 with ultrafast bootstrap and SH-aLRT (1000 replicates each) to assess the phylogenetic clades statistical support<sup>57</sup>. Trees were visualized using FigTree version 1.4.4 (http://tree.bio.ed.ac.uk/software/figtree/, accessed on 21 June 2023). Based on the clustering in the phylogenetic tree and supported by bootstrap values, sequences generated in this study are categorized into specific genotypes. Robustness of trees topology was accessed with 1000 replicates, and bootstrap values  $\geq$  70% were considered significant.

#### Deduced amino acid sequence analysis

Deduced amino acid sequences were translated with the standard genetic code using the MEGA7 software<sup>58</sup>. The GISAID RSVsurver (https://www.epicov.org/epi3/frontend#1c1efe, accessed on 04 August 2023) was used to identify key amino acid mutations. Potential N-glycosylation sites in amino acid sequences of both RSV-A and RSV-B G proteins were predicted with threshold of 0.5 by using NetNGlyc 1.0 webserver (http://www.cbs. dtu.dk/services/NetNGlyc, accessed on 04 August 2023).

#### **Statistical analysis**

SARI case-based data were entered into an EpiInfo database, merged with laboratory results, and subsequently analyzed using *R* statistical software (R.3.0.1 version). Chi-square ( $\chi^2$ ) and Fisher's exact tests were used to support the comparisons of the categorical data, where a *p*-value < 0.05 was considered statistically significant. The proportions were reported with 95% confidence intervals (CIs).

#### **Ethical considerations**

This study was conducted as part of the hospital-based surveillance of severe acute respiratory infection of the 4S (Syndromic Sentinel Surveillance in Senegal) Network, which has the approval from the Senegalese National Ethical Committee of the Ministry of Health as being less than minimal risk research. All methods were carried out in accordance with relevant guidelines and regulations, and verbal informed consent was obtained from all subjects and/or their legal guardian(s). The protocol and oral consent were determined as routine surveillance activity, and therefore non-research by the Senegalese National Ethics committee and the steering committee for 4S network, an entity representing MoH, Institut Pasteur Dakar, WHO and Clinicians in compliance with all applicable National regulations governing the protection of human subjects. Data were collected in an objective of surveillance and are anonymous. The information provided to participants was an informal description of the study. Respiratory specimens were collected, only after informed consent was granted, verbally, to local health care workers by the patients or parents in the case of minors. Patients could refuse to participate; no specimen will be taken. For the surveillance activities, written consent is judged not necessary by the Senegalese national ethics committee, which has also previously approved the work of the National Influenza Center. Collections of

non-sensitive data or an observation from normal care in which participants remain anonymous do not require ethics committee review. The patients included in this study were of all ages and consulted the Hospital sites due to acute respiratory syndromes; the patients, or parents in the case of minors, accept the tests for respiratory viruses largely because they are free and safe. The data is available in real-time to the Epidemiology Department at the Senegalese Ministry of Health and Prevention to support the appropriate public health action.

#### Data availability

All data generated during this study are contained within this manuscript. All RSV sequences generated in this study have been deposited in the GISAID database (https://gisaid.org/) under accession ID EPI\_ISL\_18228265, EPI\_ISL\_18228290, EPI\_ISL\_18228291, EPI\_ISL\_18228292, EPI\_ISL\_18228293, EPI\_ISL\_18228294, EPI\_ISL\_18228535, EPI\_ISL\_18228533, EPI\_ISL\_18228536, EPI\_ISL\_18228537, EPI\_ISL\_18228539, EPI\_ISL\_18228540, EPI\_ISL\_18228541, EPI\_ISL\_18228544, EPI\_ISL\_18228548, EPI\_ISL\_18228653, EPI\_ISL\_18228657, EPI\_ISL\_18228658, EPI\_ISL\_18228659, EPI\_ISL\_18228660, EPI\_ISL\_18228661, EPI\_ISL\_18228662, EPI\_ISL\_18228663, EPI\_ISL\_18228703, EPI\_ISL\_18228704, EPI\_ISL\_18228705, EPI\_ISL\_18228706, EPI\_ISL\_18228707, EPI\_ISL\_18228708, EPI\_ISL\_18228709, EPI\_ISL\_18228709, EPI\_ISL\_18228611, EPI\_ISL\_18229075, EPI\_ISL\_18229381, EPI\_ISL\_18229635, EPI\_ISL\_18229703.

Received: 19 September 2023; Accepted: 8 November 2023 Published online: 21 November 2023

#### References

- 1. Mazur, N. I. *et al.* The respiratory syncytial virus vaccine landscape: Lessons from the graveyard and promising candidates. *Lancet. Infect. Dis* **18**(10), e295–e311 (2018).
- Zhao, X. et al. Analysis of circulating respiratory syncytial virus A strains in Shanghai, China identified a new and increasingly prevalent lineage within the dominant ON1 genotype. Front. Microbiol. 13, 966235 (2022).
- 3. Tafa, R. B., Abebe, G. L., Tosisa, M. W. & Assefa, K. N. Molecular epidemiology of respiratory syncytial virus in children with acute respiratory illnesses in Africa: A systematic review and meta-analysis. *Journal of Global Health* **13**, 4001 (2023)
- Jarju, S. et al. Viral etiology, clinical features and antibiotic use in children < 5 years of age in The Gambia presenting with influenzalike illness. Pediatr. Infect. Dis. J. 39(10), 925–930 (2020).
- 5. Arjeyni, Y. *et al.* Molecular epidemiology of human respiratory syncytial virus in Iranian ≥ 60 years old hospitalized patients with acute respiratory symptoms. *Arch. Iran. Med.* **20**(6), 368–375 (2017).
- Chadha, M. et al. Human respiratory syncytial virus and influenza seasonality patterns—early findings from the WHO global respiratory syncytial virus surveillance. Influenza Other Respir. Viruses 14(6), 638–646 (2020).
- Shek, L. P. C. & Lee, B. W. Epidemiology and seasonality of respiratory tract virus infections in the tropics. *Paediatr. Respir. Rev.* 4(2), 105–111 (2003).
- 8. Lee, C. Y. F., Khan, S. J., Vishal, F., Alam, S. & Murtaza, S. F. Respiratory syncytial virus prevention: A new era of vaccines. *Cureus* 15(9), 45012 (2023).
- 9. Garegnani, L. et al. Palivizumab for preventing severe respiratory syncytial virus (RSV) infection in children. Cochrane Database Syst. Rev. 11 (2021).
- Garg, I., Shekhar, R., Sheikh, A. B. & Pal, S. Impact of COVID-19 on the changing patterns of respiratory syncytial virus infections. Infect. Dis. Rep. 14(4), 558–568 (2022).
- 11. Kim, H. N. *et al.* Molecular characterization of human respiratory syncytial virus in Seoul, South Korea, during 10 consecutive years, 2010–2019. *PLoS ONE* **18**(4), e0283873 (2023).
- 12. Mabilo, P., Mthiyane, H., Simane, A., Subramoney, K. & Treurnicht, F. K. Characterisation of RSV fusion proteins from South African patients with RSV disease, 2019 to 2020. *Viruses* 14(11), 2321 (2022).
- Jiang, M. L. et al. Changes in endemic patterns of respiratory syncytial virus infection in pediatric patients under the pressure of nonpharmaceutical interventions for COVID-19 in Beijing, China. J. Med. Virol. 95(1), e28411 (2023).
- 14. Fall, A. *et al.* Genetic diversity and evolutionary dynamics of respiratory syncytial virus over eleven consecutive years of surveillance in Senegal. *Infect. Genet. Evol.* **91**, 104864 (2021).
- 15. Ung, T. T. H. et al. Molecular evolution of respiratory syncytial virus subgroup A collected in north Viet Nam between 2017–2020.
- Umar, S. *et al.* Molecular epidemiology and characteristics of respiratory syncytial virus in Guangzhou, China. *World J. Pediatr.* 13, 371 (2023).
- 17. Dolores, A., Stephanie, G., Érica, G., Mistchenko, A. S. & Mariana, V. RSV reemergence in Argentina since the SARS-CoV-2 pandemic. J. Clin. Virol. 149, 105126 (2022).
- Eshaghi, A. et al. Genetic variability of human respiratory syncytial virus A strains circulating in Ontario: A novel genotype with a 72 nucleotide G gene duplication. PLoS ONE 7(3), e32807 (2012).
- Trento, A. *et al.* Major changes in the G protein of human respiratory syncytial virus isolates introduced by a duplication of 60 nucleotides. J. Gen. Virol. 84(11), 3115–3120 (2003).
- 20. Dia, N. et al. COVID-19 and Children in Senegal: Epidemiological and virological insights. Arch. Clin. Biomed. Res. 6(2), 347–357 (2022).
- 21. Dia, N. et al. COVID-19 outbreak, Senegal, 2020. Emerg. Infect. Dis. 26(11), 2771 (2020).
- Diouf, I., Bousso, A. & Sonko, I. Gestion de la pandémie COVID-19 au Sénégal. Médecine de Catastrophe-Urgences Collectives 4(3), 217–222 (2020).
- Tempia, S. et al. Decline of influenza and respiratory syncytial virus detection in facility-based surveillance during the COVID-19 pandemic, South Africa, January to October 2020. Eurosurveillance 26(29), 2001600 (2021).
- 24. Huang, Q. S. *et al.* Impact of the COVID-19 nonpharmaceutical interventions on influenza and other respiratory viral infections in New Zealand. *Nat. Commun.* **12**(1), 1001 (2021).
- 25. Adams, G. et al. The 2022 RSV surge was driven by multiple viral lineages. medRxiv 2023-01 (2023).
- Redlberger-Fritz, M., Springer, D. N., Aberle, S. W., Camp, J. V., & Aberle, J. H. The respiratory syncytial virus surge in Austria, 2022, was caused by lineages present before the COVID-19 pandemic. *bioRxiv* 2023-01 (2023).
- 27. Barry, M. A. *et al.* Performance of case definitions and clinical predictors for influenza surveillance among patients followed in a rural cohort in Senegal. *BMC Infect. Dis.* **21**(1), 1–11 (2021).
- Goya, S. et al. Toward unified molecular surveillance of RSV: A proposal for genotype definition. Influenza Other Respir. Viruses 14(3), 274–285 (2020).
- Fall, A. et al. Epidemiology and molecular characterization of human respiratory syncytial virus in Senegal after four consecutive years of surveillance, 2012–2015. PLoS ONE 11(6), e0157163 (2016).

- Munkstrup, C. et al. Early and intense epidemic of respiratory syncytial virus (RSV) in Denmark, August to December 2022. Eurosurveillance 28(1), 2200937 (2023).
- Delestrain, C. et al. Impact of COVID-19 social distancing on viral infection in France: A delayed outbreak of RSV. Pediatr. Pulmonol. 56(12), 3669–3673 (2021).
- Opek, M. W. et al. Delayed respiratory syncytial virus epidemic in children after relaxation of COVID-19 physical distancing measures, Ashdod, Israel, 2021. Eurosurveillance 26(29), 2100706 (2021).
- 33. Reicherz, F. et al. Waning immunity against respiratory syncytial virus during the coronavirus disease 2019 pandemic. J. Infect. Dis. 226(12), 2064–2068 (2022).
- Malasao, R. et al. Predominance of ON1 and BA9 genotypes of human respiratory syncytial virus in children with acute respiratory infection in Chiang Mai, Thailand, 2020–2021. J. Infect. Public Health 16(9), 1418–1426 (2023).
- Tabatabai, J. et al. Molecular epidemiology of respiratory syncytial virus in hospitalised children in Heidelberg, Southern Germany, 2014–2017. Infect. Genet. Evol. 98, 105209 (2022).
- 36. Lanari, M. *et al.* Prevalence of respiratory syncytial virus infection in Italian infants hospitalized for acute lower respiratory tract infections, and association between respiratory syncytial virus infection risk factors and disease severity. *Pediatr. Pulmonol.* **33**(6), 458–465 (2002).
- Korsun, N. et al. Predominance of ON1 and BA9 genotypes of respiratory syncytial virus (RSV) in Bulgaria, 2016–2018. J. Med. Virol. 93(6), 3401–3411 (2021).
- 38. Xie, Z. *et al.* The burden of respiratory syncytial virus associated with acute lower respiratory tract infections in Chinese children: A meta-analysis. *Transl. Pediatr.* **9**(4), 496 (2020).
- Pierangeli, A. et al. Genetic diversity and its impact on disease severity in respiratory syncytial virus subtype-A and-B bronchiolitis before and after pandemic restrictions in Rome. J. Infect. 87, 305–314 (2023).
- Lin, T. Y. et al. Outbreak of respiratory syncytial virus subtype ON1 among children during COVID-19 pandemic in Southern Taiwan. J. Microbiol. Immunol. Infect. 55(6), 1168–1179 (2022).
- 41. Hall, C. B. et al. The burden of respiratory syncytial virus infection in young children. N. Engl. J. Med. 360(6), 588-598 (2009).
- 42. Lin, H. C. et al. RSV pneumonia with or without bacterial co-infection among healthy children. J. Formosan Med. Assoc. 121(3), 687–693 (2022).
- 43. Harada, Y. *et al.* Does respiratory virus coinfection increase the clinical severity of acute respiratory infection among children infected with respiratory syncytial virus?. *Pediatr. Infect. Dis. J.* **32**(5), 441–445 (2013).
- 44. Luo, H. J. *et al.* Epidemiological characteristics and phylogenic analysis of human respiratory syncytial virus in patients with respiratory infections during 2011–2016 in southern China. *Int. J. Infect. Dis.* **90**, 5–17 (2020).
- Heppe-Montero, M., Walter, S., Hernández-Barrera, V., Gil-Prieto, R. & Gil-de-Miguel, Á. Burden of respiratory syncytial virusassociated lower respiratory infections in children in Spain from 2012 to 2018. BMC Infect. Dis. 22(1), 315 (2022).
- González-Ortiz, A. M. et al. Severe respiratory syncytial virus infection in hospitalized children. Arch. Med. Res. 50(6), 377–383 (2019).
- Tabor, D. E. et al. Global molecular epidemiology of respiratory syncytial virus from the 2017–2018 INFORM-RSV Study. J. Clin. Microbiol. 59(1), 10–1128 (2020).
- Kobiałka, M., Jackowska, T. & Wrotek, A. Risk factors for severe respiratory syncytial virus infection in hospitalized children. Viruses 15(8), 1713 (2023).
- Greenberg, D., Dagan, R., Shany, E., Bar-Ziv, J. & Givon-Lavi, N. Increased risk for respiratory syncytial virus-associated, community-acquired alveolar pneumonia in infants born at 31–36 weeks of gestation. *Pediatr. Infect. Dis. J.* 33(4), 381–386 (2014).
- Goya, S. et al. Genomic characterization of respiratory syncytial virus during 2022–23 outbreak, Washington, USA. Emerg. Infect. Dis. 29(4), 865 (2023).
- Chen, X. et al. A multi-center study on molecular epidemiology of human respiratory syncytial virus from children with acute lower respiratory tract infections in the mainland of China between 2015 and 2019. Virol. Sin. 36(6), 1475–1483 (2021).
- 52. Fitzner, J. et al. Revision of clinical case definitions: Influenza-like illness and severe acute respiratory infection. Bull. World Health Organ. 96(2), 122 (2018).
- Walker, G. et al. Genomic and immunological characterisation shows increased respiratory syncytial virus cases not due to waning antibody mediated immunity. SSRN 4515871 (2023).
- 54. Hale, C. Twist Bioscience launches research panel for screening COVID-19, other respiratory viruses. Fiercebiotech.com (2020).
- Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. Mol. Biol. Evol. 30(4), 772–780 (2013).
- Hall, T. A. BioEdit: A user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. Nucleic Acids Symp. Ser. 41(41), 95–98 (1999).
- 57. Minh, B. Q. et al. IQ-TREE 2: New models and efficient methods for phylogenetic inference in the genomic era. *Mol. Biol. Evol.* **37**(5), 1530–1534 (2020).
- Kumar, S., Stecher, G., & Tamura, K. MEGA7: Molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Mol. Biol. Evol.* 33(7), 1870–1874 (2016).

# Acknowledgements

We would like to acknowledge the Ministry of Health for its support and all the SARI sentinel sites healthcare workers. We convey special thanks to Catherine Commaille-Chapus and Camille Hunsinger from Impact Healthcare for the coordination and quality data verifications, and for their unwavering support.

# **Author contributions**

N.D., M.A.B., F.T., J.B.N.D., D.B. and G.F. conceptualized the project; M.M.J., M.M.D., M.P.M., N.K.N., D.K. and S.S. performed the experiments; F.T., J.B.N.D., D.B., M.M.J., S.S., D.G., M.A.B., C.L. and N.D. collected and analyzed the data; M.M.J. wrote the original draft preparation; N.D. and G.F. reviewed and edited the manuscript. All authors read and approved the final manuscript.

# Funding

This study was conducted in the purpose of both the global influenza hospital surveillance network (GISHN) program and the Africa CDC Pathogen Genomics Initiative funds (CARES grant 4306–22-EIPHLSS-GENOMICS).

# **Competing interests**

The authors declare no competing interests.

# Additional information

**Correspondence** and requests for materials should be addressed to N.D.

Reprints and permissions information is available at www.nature.com/reprints.

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

© The Author(s) 2023